

LIBRARY

NOV 22 1999

LIBRARY
UNIVERSITY

First record of *Metacyclops stammeri* Kiefer, 1938 from Balkan Peninsula

(Crustacea, Copepoda)

Tomislav Karanovic

Karanovic, T. (1999): First record of *Metacyclops stammeri* Kiefer, 1938 from Balkan Peninsula (Crustacea, Copepoda). – *Spixiana* **22/3**: 193-198

The cyclopoid copepod *Metacyclops stammeri* Kiefer, 1938 (Copepoda, Cyclopoidea) is recorded from Balkan Peninsula for the first time. This species was known only from the Apulian region in the southern part of Italy. A detailed description of that interesting Tertiary relict is given.

Tomislav Karanovic, Institute of Marine Biology, P.O. Box 69, 85335 Kotor, Montenegro, Yugoslavia

Introduction

Metacyclops was first established as a subgenus of the genus *Mesocyclops* by Kiefer (1927, 1928), and later as a genus (Kiefer 1937, 1960). Lindberg (1961) redefined the genus *Metacyclops* and also gave a key to females. He emphasized the distinctive features of that genus. One unsuccessful attempt to unite the three genera *Apocyclops*, *Psammophilocyclops*, and *Metacyclops* as subgenera of the genus *Metacyclops* was made by Plesa (1981). This grouping was followed by Pesce (1985a), but Reid (1987) showed the unnecessary of that attempt. Forty-seven currently recognized taxa were listed in a world key by Herbst (1988), together with their geographical distribution. Reid (1991) added to the above list *Mesocyclops pseudoanceps*, which proved to be a *Metacyclops*. Recently described species and subspecies not included in Herbst's key are: *M. postojnae* Brancelj, 1987 from Slovenia; *M. janstocki* Herbst, 1990 from Antigua; *M. leptosus totaensis* Reid et al. 1990 from Colombia; *M. cushae* Reid, 1991 from the United States; *M. oraemaris* Rocha, 1994 from Brazil; *M. hirsutus* Rocha, 1994 also from Brazil; *M. mendocinus insulensis* Defaye & Dussart, 1991 from the Azores; and *M. geltrudeae* Galassi & Pesce, 1994 from Venezuela. Therefore, now the genus *Metacyclops* includes fifty-six species and subspecies of which the majority are tropical and temperate in distribution.

Kiefer (1938) described two species of the genus *Metacyclops* from the Apulian region in Italy: *M. subdohus* and *M. stammeri*. *M. subdohus* was latter rediscovered in Apulia (Pesce et al. 1978), and was also found on the Balears, on Corsica and in Greece (Lindberg 1956, Pesce 1978, 1983, Pesce & Maggi 1981, 1983). *M. stammeri* was also rediscovered in several localities in Apulia (Pesce et al. 1978), but was never found outside that region. During an investigation of the copepod fauna in Montenegro two males of *Metacyclops stammeri* were found. It is the first record of that species from Balkan Peninsula.

Methods

Samples were collected with different types of hand-nets and little rubber pumps. The material was preserved by adding several drops of 36% formaldehyde, and very soon after that washed, and copepods were separated and removed into 70% ethanol. One specimen was dissected in a mixture of distilled water and glycerol (1:1),

with fine entomological needles. All drawings have been prepared using a drawing attachment on a Leica DMLS microscope with C-PLAN achromat objectives. Dissected appendages were preserved in Faure's medium. The not dissected specimen was, after examination, again preserved into 70 % ethanol.

Abbreviations used in the text and figures: Fu: furca; A1: antennula; A2: antenna; Mx: maxilla; P1-P6: 1st-6th leg; Enp: endopodite; Exp: exopodite; Enp2P4: second endopodite article of the 4th leg.

Results

Metacyclops stammeri Kiefer, 1938

Metacyclops Stammeri Kiefer, 1938, p. 7, figs 11-17;

Metacyclops stammeri, Lindberg 1961, p. 142; Dussart 1969, p. 173, fig. 85; Pesce et al. 1978, p. 37, fig. 5; Kiefer 1978, p. 215; Pesce 1985, p. 135; Dussart & Defaye 1985, p. 100; Lescher-Moutoué 1986, p. 308; Herbst 1988, p. 150.

Material examined: Two ♂♂ from the cave Sutimska Jama (42°25'50" N, 19°10'40" E), near the town Podgorica, Montenegro, collected by T. Karanovic, 3 February 1997. One ♂ was completely dissected and mounted on a slide in Faure's medium (No. 8/66/0584/i). The other ♂ is preserved in 70 % ethanol in a glass test tube (No. 7/0584/i). Both specimens are deposited in the author's working collection, at the Institute of Marine Biology, Kotor, Montenegro.


Description

Male. Length of dissected specimen, including furcal rami (excluded furcal setae) 0.54 mm. Length of non-dissected specimen 0.593 mm. Body surface without patterns of cuticle striae or setulose areas, and colourless. Nauplius eye absent. Hind margin of each body segment smooth and without rows of spines (Fig. 6). Cephalothorax very long (longer than thoracic somites and four abdominal somites together). Anal somite also without rows of spines at the base of furcal rami, both ventrally and dorsally (Figs 1, 2). Anal operculum very short and broad. Furcal rami parallel, close, and about 4 times longer than wide (Fig. 1). Lateral seta inserted at distal three-fifths of ramus. Apical setae with following proportion of lengths (from outside to inside: 1/2.8/8.4/1. Inner-middle seta (the longest one) swollen at base, and without cuticular pint. Dorsal seta attached almost at the end of the furcal ramus, and about 1.4 times longer than it. That seta is also longer than outer-middle seta (Fig. 1). A1 geniculate, 14-segmented, and with aesthetascs only at first and third segments (Fig. 3). A2 4-segmented, with well-developed exopodite seta, and without rows of spines or hairs on the first segment (Fig. 10). All swimming legs with 2-segmented exopodites and endopodites. Spine formula of distal exopodite articles P1-P4: 3.4.4.3; setal formula of those articles: 5.5.5.5 (Figs 7, 9, 11, 12). Enp2P4 approximately 3.2 times as long as width, apically armed with outer spine and inner plumose seta (which is more than twice longer than spine). Intercoxal plates in all swimming legs smooth and without hairs or spinules. P5 consisting of 1 fused and 1 free article (Fig. 4). Free article longer than width, armed apically with very long outer plumose seta, and inner spine (which is about 1.3 times longer than article that bears it). Seta on free article 1.5 times longer than seta on fused article. P6 armed with two appendages: inner spine and outer seta (Fig. 8).


Discussion

Kiefer (1938) described *M. stammeri* from three caves (La Zinzuleisa, Abissa, and Grotta dei Diavoli), south from the town Otranto, in the Apulian region of southern Italy. Pesce et al. (1978) rediscovered it in the same caves, and also on four additional localities near the town Gallipoli, also in the Apulian region (Fig. 13), although their geographical coordinates are erroneous. Dussart & Defaye (1985) by mistake recorded the species from Corsica, Sicily, and Sardinia, as well as Herbst (1988), probably because of the wrong interpretation of Kiefer (1978). Specimens from Montenegro are very similar to those from the Apulian region, and there is no doubt of their specific status. They differ just by a little longer dorsal setae on Fu.

Metacyclops stammeri Kiefer, 1938 is clearly distinguishable from all species in the genus by the shape of Enp2P4 and Fu. Only *M. postojnae* Brancelj (1987) from Slovenia possesses similar Enp2P4, but it has different body habitus and shape of Fu. Similar body habitus and reduction of outer-middle seta



Figs 1-6. *Metacyclops stammeri* Kiefer, 1938. 1-5. ♂ (0.54 mm). 6. ♂ (0.593 mm). 1. Anal somite & Fu, dorsal view. 2. Abdomen, ventral view. 3. A1. 4. P5. 5. Mx. 6. Habitus, dorsal view. Scales: 0.1 mm.



Figs 7-12. *Metacyclops stammeri* Kiefer, 1938, ♂ (0.54 mm). 7. P4. 8. P6. 9. P3. 10. A2. 11. P1. 12. P2. Scales: 0.1 mm.




Fig. 13. Distribution of *Metacyclops stammeri* Kiefer, 1938 (circles: localities; broken line: supposed border of distribution).

on Fu has *M. trisetosus* (Herbst, 1957), described from Croatia and rediscovered in northern Italy by Stoch (1988), but it differs by the shape of P4, and the outer-middle seta on Fu is completely reduced in that species.

M. stammeri is one of the Tertiary relicts that have survived Quaternary cyclic periods of the ice ages in the subterranean waters. Its recent distribution (Fig. 13) is probably the consequence of the last ice age, which drive it into two refuge pockets (Apulian karstic peninsula, and Skadar karstic valley). Its return to the north areas in Balkan Peninsula is thwart by geographical obstacles and also by the presence of three subterranean species from the genus *Metacyclops* in central and northern part of the Dinaric Alps: *M. trisetosus* (Herbst, 1957), *M. postojnae* Brancelj, 1987, and *M. gasparoi* Stoch, 1987. In Italy the most important obstacle probably is the distribution of main karstic areas (see Pesce 1985: fig. 1).

References

- Brancelj, A. 1987. Cyclopoida and Calanoida (Crustacea, Copepoda) from the Postojna-Planina Cave System (Slovenia). – Biol. Vestn. Ljubljana 35: 1-16
- Defaye, D. & B. Dussart. 1991. Sur les Crustacés Copepodes des Açores. – Anns Limnol. 27(2): 119-132
- Dussart, B. 1969. Les Copepodes des eaux continentales d'Europe Occidentale, II. Cyclopoïdes et Biologie. – Edit. Boubee & Cie, Paris: 292 pp.

- & D. Defaye. 1985. Repertoire Mondial des Copepodes Cyclopoïdes. – Edit. du C.N.R.S.: 236 pp.
- Galassi, D. P. & G. L. Pesce. 1984. *Metacyclops geltrudeae* n. sp., a new cyclopoid from ground waters of Venezuela (Copepoda, Cyclopidae). – Crustaceana 67(3): 284-287
- Herbst, H.-V. 1957. Zoological Results of a Collecting Journey to Yugoslavia, 1954, 5. Cyclopoida Gnathostoma (Crustacea Copepoda). – Beaufortia 65(5): 223-240
- 1988. Zwei neue *Metacyclops* (Crustacea) von den Westindischen Inseln Barbados und Aruba: *M. agnitus* n. sp. und *M. mutatus* n. sp., sowie ein Bestimmungsschlüssel für das Genus. – Bijdr. Dierk. 58(1): 137-154
- 1990. *Metacyclops janstocki* n. sp. (Crustacea, Copepoda) von Antigua (Westindische Inseln). – Beaufortia 41: 75-81
- Kiefer, F. 1927. Versuch eines Systems der Cyclopiden. – Zool. Anz. 73(11/12): 302-308
- 1928. Über Morphologie und Systematik der Süßwasser-Cyclopiden. – Zool. Jahrb. Syst. 54(5/6): 495-556
- 1937. Über Systematik und geographische Verbreitung einiger Gruppen stark verkümmelter Cyclopiden (Crustacea Copepoda). – Zool. Jahrb. Syst. 70(5/6): 421-442
- 1938. Cyclopiden (Crustac. Cop.) aus süditalienischen Brunnen und Höhlen. – Zool. Anz. 123(1/2): 1-12
- 1960. Ruderfusskrebse (Copepoden). – Kosmos Verlag Franckh, Stuttgart: 97 pp.
- 1978. Copepoda non-parasitica. – In: Limnofauna Europaea, ed. J. Illies, Gustav Fischer Verlag: 209-225
- Lescher-Moutoué, F. 1986. Copepoda Cyclopoida Cyclopidae des eaux douces souterraines continentales. – In: Stygofauna Mundi, ed. L. Botosaneanu: 299-312
- Lindberg, K. 1956. Cyclopides (Crustaces Copepodes) de la Sardaigne. – Mem. Soc. Entomol. Ital., Genova 35: 71-79
- 1961. Remarques sur le genre *Metacyclops* (Kiefer 1927) et description d'un *Metacyclops* nouveau du Portugal. – Kungl. Fysiogr. Sällsk. Lund Forh. 31(14): 133-145
- Pesce, G. L. 1978. The occurrence of *Metacyclops subdolosus* Kiefer (Crustacea: Copepoda) in subterranean waters of Greece with remarks on its systematic status. – Int. J. Speleol. 10: 179-183
- 1983. Some remarks on the stygofauna of Greece. – Biologia Gallo-Hellenica 10: 103-112
- 1985. The groundwater fauna of Italy: a synthesis. – Stygologia 1(2): 129-159
- 1985a. Cyclopids (Crustacea, Copepoda) from West Indian groundwater habitats. – Bijdr. Dierk. 55(2): 295-323
- , G. Fusacchia, D. Maggi & P. Tete. 1978. Ricerche faunistiche in acque freatiche del Salento. – Thalassia Salentina 8: 1-51
- & D. Maggi. 1981. Cyclopides et calanoides des eaux phreatiques de la Grece meridionale et insulaire (Crustacea: Copepoda). – Ecologia Mediterranea 7: 163-182
- & --. 1983. Ricerche faunistiche in acque sotterranee freatiche della Grecia Meridionale ed Insulare e stato attuale delle conoscenze sulla stigofauna di Grecia. – Natura Soc. Ital. Sci. Nat. Mus. Civ. Stor. Nat. Acquario, Milano 74(1/2): 15-73
- Plesa, C. 1981. Cyclopides (Crustacea, Copepoda) de Cuba. – Result. exp. biospeol. cubano-roumaines a Cuba 3: 17-34
- Reid, J. W. 1987. The cyclopoid copepods of a wet campo marsh in central Brazil. – Hydrobiologia 153: 121-138
- 1991. The genus *Metacyclops* (Copepoda: Cyclopoida) present in North America: *M. cushae* new species, from Louisiana. – Jour. Crustac. Biol. 11(4): 639-646
- , J. A. M. Arevalo & M. M. Fukushima. 1990. *Metacyclops leptosus totaensis*, new subspecies (Crustacea: Copepoda) from Lago de Tota, Colombia. – Proc. Biol. Soc. Washington 103: 674-680
- Rocha, C. E. F. 1994. New species of *Metacyclops* (Copepoda, Cyclopidae) from Brazil, with remarks on *M. campestris*. – Zoologica Scripta 23(2): 133-146
- Stoch, F. 1987. Cave-dwelling cyclopoïdes (Crustacea, Copepoda) from Venezia Giulia (northeastern Italy). – Bull. Zool. Mus. Univ. Amsterdam 11(5): 41-55
- 1988. Secondo contributo alla conoscenza dei Calanoidi e dei Ciclopoidi (Crustacea, Copepoda) delle acque carsiche sotterranee della Venezia Giulia (Italia Nordorientale). – Atti Mem. Comm. Grotte "E. Boegan" 27: 63-71