


Proceedings of  
the United States  
National Museum



SMITHSONIAN INSTITUTION • WASHINGTON, D.C.

Volume 124

1967

Number 3628

Cyclopoid Copepods of the Genus *Paranthesius*  
Associated with Marine Pelecypods in Chile

By Arthur G. Humes<sup>1</sup>

Records of the genus *Paranthesius* from the western coast of North and South America are relatively scarce. Thompson (1897, p. 87) in Herdman, Thompson, and A. Scott, described *Pseudolichomolgus columbiae* (= *Paranthesius columbiae* according to Illg, 1949) from a single planktonic specimen in Puget Sound, Washington. Illg (1949) reported five species of *Paranthesius* from various pelecypods in California: *P. columbiae* (Thompson, 1897) from *Schizothaerus nuttalli* (Conrad) and *Protothaca tenerrima* (Carpenter), and four new species, *P. panopeae* from *Panope generosa* Gould, *P. tivelae* from *Tivela stultorum* (Mawe), and *P. saxidomi* and *P. perplexus* from *Saxidomus nuttalli* Conrad.

On Cruise 18 of the R/V *Anton Bruun*, during the Southeastern Pacific Biological Oceanographic Program of the National Science Foundation, the ship visited Valparaiso and Iquique, Chile, where the author collected specimens of the two new copepods described below from pelecypods purchased in the local fish markets. The first of the new species is the second record of *Paranthesius* from the genus *Protothaca* and the second represents the first record from *Mesodesma*.

<sup>1</sup> Department of Biology, Boston University, Boston, Mass.

The discovery of these two new copepods extends the eastern Pacific range of the genus as far south as central Chile.

The study of the specimens has been aided by a grant (GB-5838) from the National Science Foundation.

All figures have been drawn with the aid of a camera lucida. The letter after the explanation of each figure refers to the scale at which it was drawn. The abbreviations used are:  $A_1$ =first antenna,  $A_2$ =second antenna, L=labrum, MD=mandible, P=paragnath,  $MX_1$ =first maxilla,  $MX_2$ =second maxilla, MXP= maxilliped, and  $P_1$ =leg 1.

I thank Prof. José Stuardo of the Universidad de Concepción, Concepción, Chile, for the identifications of the two pelecypod hosts.

## Family LICHOMOLGIDAE Kossmann, 1877

### Genus *Paranthesius* Claus, 1889

#### *Paranthesius protothacae*, new species

#### FIGURES 1-33

TYPE MATERIAL.—11 ♀♀ and 10 ♂♂ from the mantle cavity of 59 *Protothaca thaca* Molina purchased in the fish market at Iquique, Chile, Aug. 23, 1966. (These pelecypods probably came from Punta Guanillos, 21°15'S, 70°07'W, south of Iquique.) Holotype female, allotype, and 16 paratypes (8 ♀♀, 8 ♂♂) deposited in the United States National Museum, and the remaining paratypes (dissected) in the collection of the author.

FEMALE.—Body (figs. 1, 2) moderately slender, about three times longer than wide, with prosome not greatly thickened dorsoventrally. Length (not including setae on caudal rami) 1.62 mm (1.47–1.78 mm) and greatest width 0.54 mm (0.50–0.61 mm), based on 10 specimens measured in lactic acid. Dorsoventral thickness of prosome about 0.45 mm. Ratio of length to width of prosome 1.9:1. Segment of leg 1 separated from head dorsally and laterally by a distinct furrow. Epimeral areas of metasomal segments as illustrated.

Segment of leg 5 (fig. 3)  $91\mu \times 135\mu$ . Between this segment and genital segment a weak ventral intersegmental sclerite, its sclerotization more evident laterally (fig. 4). A median prominence in front of segment of leg 5 (figs. 4, 5). Genital segment as long as wide,  $216\mu \times 216\mu$ , in dorsal view anterior half laterally expanded with rounded margins and posterior half narrower and tapering slightly posteriorly with straight margins. Areas of attachment of egg sacs located dorso-laterally, each area (fig. 6) bearing small seta  $13\mu$  long and spiniform process  $6\mu$ . Three postgenital segments  $78\mu \times 120\mu$ ,  $70\mu \times 109\mu$ , and  $114\mu \times 99\mu$ , from anterior to posterior. Anal segment with postero-ventral margins unornamented.

Caudal ramus (fig. 7) elongated,  $135\mu \times 34\mu$  in greatest dimensions, four times longer than wide. Outer lateral seta  $52\mu$ , outermost terminal seta  $70\mu$ , innermost terminal seta  $57\mu$ . Two long median terminal setae  $195\mu$  (outer) and  $380\mu$  (inner) and inserted slightly dorsally. Dorsal pedicellate seta small,  $18\mu$  long. All setae naked. Dorsal surface of ramus without ornamentation.

Dorsal surface of prosome and urosome with a few hairs (as in fig. 1); ventral surface of urosome almost devoid of ornamentation. Ratio of length of prosome to that of urosome 1.4:1.

Egg sac unknown.

Rostrum (figs. 8, 9) a rounded prominence with a rather blunt posteroventral tip.

First antenna (fig. 10) slender (about  $358\mu$  in length) and 7-segmented, with a sclerite on third segment (fig. 11) suggesting an intercalary segment. Lengths of segments (measured along posterior nonsetiferous margins)  $13\mu$  ( $50\mu$  along anterior margin),  $81\mu$ ,  $35\mu$ ,  $60\mu$ ,  $55\mu$ ,  $41\mu$ , and  $36\mu$  respectively. Formula for armature, as in many other lichomolgids, 4, 13, 6, 3, 4+1 aesthete, 2+1 aesthete, and 7+1 aesthete. All setae naked.

Second antenna (fig. 12) 4-segmented. Armature 1, 1, 3, and I,5. All setae naked. Three long terminal setae on last segment jointed; claw pale brown,  $77\mu$  along its axis. Second segment with few short spinules along outer edge.

Labrum (figs. 13, 18) with two widely divergent posteroventral lobes.

Mandible (fig. 14) with a slender elongated blade bearing spinules along each side. Paragnath (fig. 13) a small hairy lobe. First maxilla (fig. 15) with four setae (in one first maxilla of one female only three, seta nearest to two terminal ones being absent). Second maxilla (fig. 16) 2-segmented. First segment unornamented. Second segment having on outer (ventral) margin a small proximal seta, bearing on inner surface a seta with very short barbules and a long seta with spinules along one edge, and terminating in a long lash with long spinules on one margin and few small spinules on opposite side. Maxilliped (fig. 17) 3-segmented. First segment lacking spines or setae. Second segment with two rather unequal setae. Small third segment terminating in a spiniform process (not clearly articulated with segment) with a small seta near base. Postero-outer surface of second segment and postero-outer distal part of first segment with dense covering of fine short hairlike ornamentation.

Postoral area as in figure 18. Median region between mandibles, paragnaths, and first maxillae slightly protuberant. A sclerotized line between bases of maxillipeds.

Legs 1-4 (figs. 19-22) with trimerous rami. Armature as follows (Roman numerals indicating spines, Arabic numerals setae):

P <sub>1</sub>	protopod	0-1	1-0	exp	I-0	I-1	III,I,4
				end	0-1	0-1	I,5
P <sub>2</sub>	protopod	0-1	1-0	exp	I-0	I-1	III,I,5
				end	0-1	0-2	I,II,3
P <sub>3</sub>	protopod	0-1	1-0	exp	I-0	I-1	III,I,5
				end	0-1	0-2	I,II,2
P <sub>4</sub>	protopod	0-1	1-0	exp	I-0	I-1	II,I,5
				end	0-1	0-1	II

Inner margin of basis ornamented with spinules in leg 1, with hairs in legs 2-4. Leg 4 with last segment of exopod most often with the formula II,I,5 (in seven females), but occasionally III,I,5 (in two females) or II,I,5 and III,I,5 (in two females where formula differed on opposite legs). Last segment of endopod in leg 4 measuring  $74\mu$  x  $28\mu$  in greatest dimensions, with two terminal fringed spines  $64\mu$  (outer) and  $79\mu$  (inner).

Leg 5 (fig. 23) with its free segment  $52\mu$  x  $19\mu$ ; ornamented with few minute distal outer spinules and armed with a terminal naked seta  $77\mu$  and a subterminal inner lamellate spine  $33\mu$  with blunt tip. Seta on body near insertion of free segment  $36\mu$  and naked.

Leg 6 probably represented by small seta and spiniform process near attachment of egg sac (see fig. 6).

Color in life in reflected light transparent to slightly amber, eye red.

MALE.—Body (figs. 24, 25) slender as in the female. Length (without setae on the caudal rami) 1.44 mm (1.22-1.54 mm) and greatest width 0.44 mm (0.38-0.50 mm), based on 10 specimens measured in lactic acid. Ratio of length to width of prosome 1.8:1.

Segment of leg 5 (fig. 26)  $52\mu$  x  $109\mu$ . Between this segment and genital segment no ventral intersegmental sclerite. Genital segment only slightly longer than wide,  $255\mu$  x  $234\mu$ , with lateral margins gently rounded in dorsal view. Four postgenital segments  $52\mu$  x  $99\mu$ ,  $55\mu$  x  $94\mu$ ,  $49\mu$  x  $84\mu$ , and  $88\mu$  x  $81\mu$ , from anterior to posterior.

Caudal ramus resembling that of female,  $114\mu$  x  $29\mu$ .

Surfaces of body ornamented with few hairs (fig. 24) as in female. Ratio of length of prosome to that of urosome 1.2:1.

Rostrum like that of female.

First antenna (fig. 27) resembling that of female, but with three aesthetes added, so that the formula is 4, 13 + 2 aesthetes, 6, 3 + 1 aesthete, 4 + 1 aesthete, 2 + 1 aesthete, and 7 + 1 aesthete. Second antenna (fig. 28) similar to that of female, but with four elements on third segment and with small spinules along inner edge of second segment.

Labrum, mandible, paragnath, first maxilla, and second maxilla as in female. Maxilliped (fig. 29) slender and 4-segmented, assuming that proximal part of claw represents a fourth segment. First segment unarmed. Second segment bearing two inner setae, innermost with sclerotized basal portions (fig. 30), and row of spinules on posterior surface. Third segment small and unarmed. Recurved claw  $172\mu$  along axis, with slight indication of division about midway; terminal lamella very narrow; two unequal setae on proximal part of claw.

Postoral area as in female.

Legs 1-4 as in female, with same spine and setal formula. Last segment of endopod of leg 1 (fig. 31) showing very slight sexual dimorphism, with more numerous terminal spinules and outer fringe on spine more strongly spinulose than in female.

Leg 5 (fig. 32) with free segment  $33\mu \times 13\mu$ , terminal seta  $62\mu$ , and subterminal inner spine  $15\mu$  and without lamellae.

Leg 6 (fig. 33) a posteroventral flap on genital segment bearing two naked setae  $29\mu$  and  $38\mu$  long, with a row of minute spinules near insertion of longer more anterior seta.

Spermatophore not seen.

Color in life like that of female.

ETYMOLOGY.—The specific name *protothacae* is formed from the generic name of the host.

COMPARISON WITH RELATED SPECIES.—Following the views of Illg (1949) and adding information from Bocquet and Stock (1958, 1958a, 1959), Gotto (1961) regarded the genus *Paranthesius* as embracing 21 species. Stock (1964) has, however, removed six of these species to the genus *Scambicornus* Heegaard, 1944. These are *S. prehensilis* (Sars, 1918), *S. robustus* (Thompson and A. Scott, 1903), *S. serendibicus* (Thompson and A. Scott, 1903), *S. propinquus* (Nicholls, 1944), *S. finmarchicus* (T. Scott, 1903), and *S. tenuicaudis* (Sars, 1918). A recently described form, *P. colmani* Reddiah, 1960, must be added to the genus, bringing the number of currently recognized species to 16. For the purpose of the description of *P. protothacae* and the following new species, *Paranthesius* is held to consist of these 16 species, although it is recognized that many problems exist in the determination of the generic limits and that several authors (for example, Monod and Dollfus, 1932; Illg, 1949; Sewell, 1949; Bocquet and Stock, 1957; and Reddiah and Williamson, 1959) have variously interpreted the scope of the genus.

Nine species of *Paranthesius* have a distinct beak on the rostrum and may thus be distinguished readily from *P. protothacae*. These are: *P. columbiae* (Thompson, 1897); *P. panopeae* Illg, 1949; *P. tivelae* Illg, 1949; *P. nasutus* (Edwards, 1891); *P. pectinis* (Pesta,

1909); *P. rostratus* (Canu, 1891); *P. haploceras* Bocquet and Stock, 1959; *P. barneae* (Pelseneer, 1929); and *P. colmani* Reddiah, 1960.

The new species may be separated from the remaining seven species as follows: *P. anemoniae* Claus, 1889, has an elongated segment in leg 5 (about 5:1 in Bocquet and Stock, 1959a, fig. 5d) and the second antenna has three terminal claws; *P. cynthiae* (Brian, 1924) has the caudal ramus equal to the length of the last two postgenital segments and the last segment of the second antenna much longer than the penultimate segment; *P. myxicolae* Bocquet and Stock, 1958, has recurved spines on the first and second segments of the first antenna and three terminal claws on the second antenna; *P. parvus* (Norman and T. Scott, 1905) has jointed setae (no claw) on the last segment of the second antenna in the female; *P. perplexus* Illg, 1949, has an ellipsoid rounded process on each side of the genital segment in the female; *P. saxidomi* Illg, 1949, has a reduced maxilliped in the female (length less than half the basal segment of the second maxilla), the basis of legs 1-3 with stout spinules, the last segment of the endopod of leg 4 with the inner spine nearly 2.5 times the length of the outer spine, the segment of leg 5 in the female with a rounded expansion at the base and the two terminal elements subequal in length, the male maxilliped with a modified seta on the second segment, and the male genital segment with two pairs of rows of spinules; and *P. validus* (Sars, 1918) has a caudal ramus that in the female is only slightly longer than wide.

*Paranthesius mesodesmatis*, new species

FIGURES 34-43

TYPE MATERIAL.—14 ♀♀ and 16 ♂♂ from the mantle cavity of 39 *Mesodesma donacium* Lamarck purchased in the fish market at Valparaiso, Chile, Aug. 4, 1966. (These pelecypods presumably came from Viña del Mar, near Valparaiso.) Holotype female, allotype, and 24 paratypes (11 ♀♀, 13 ♂♂) deposited in the United States National Museum, and the remaining paratypes (dissected) in the collection of the author.

In the following description those features not mentioned may be assumed to be essentially like those of the previous species.

FEMALE.—Body (figs. 34, 35) with tumid prosome. Length (not including setae on caudal rami) 1.49 mm (1.40-1.63 mm) and greatest width 0.64 mm (0.48-0.77 mm), based on 8 specimens measured in lactic acid. Dorsoventral thickness of prosome about 0.66 mm in a specimen 1.50 x 0.64 mm. (One female less tumid than the others, its dimensions 1.47 x 0.47 mm, with dorsoventral thickness of 0.42 mm.) Ratio of length to width of prosome 1.38:1. Segment of leg 1 very

weakly delimited from head. Epimeral areas of first and second metasomal segments apparently distorted by swelling of prosome (fig. 34).

Segment of leg 5 (fig. 36)  $69\mu \times 133\mu$ . Genital segment  $195\mu \times 190\mu$ . Areas of attachment of egg sacs dorsolateral. Three postgenital segments  $73\mu \times 104\mu$ ,  $57\mu \times 91\mu$ , and  $91\mu \times 86\mu$ , from anterior to posterior. Caudal ramus  $102\mu \times 30\mu$ , 3.4 times longer than wide, slightly shorter than in *P. protothacae*.

Ratio of length of prosome to that of urosome 1.5:1.

Egg sac (fig. 34) elongated,  $1.25 \times 0.26$  mm in one female,  $1.73 \times 0.29$  mm in another, with many eggs each about  $86\mu$  in diameter.

Rostrum (fig. 37) as in previous species.

First antenna segmented and armed as in *P. protothacae*, but shorter, about  $285\mu$  long. Lengths of segments (measured along posterior non-setiferous margins)  $16\mu$  ( $42\mu$  along anterior margin),  $61\mu$ ,  $25\mu$ ,  $51\mu$ ,  $43\mu$ ,  $36\mu$ , and  $27\mu$  respectively. All setae naked.

Second antenna segmented and armed as in *P. protothacae*. Claw  $59\mu$  along axis.

Labrum, mandible, paragnath, first maxilla, and second maxilla as in *P. protothacae*. Maxilliped resembling that of previous species, but the two setae on second segment more nearly equal and fine hairlike ornamentation on first two segments apparently absent.

Postoral area as in previous species.

Legs 1–4 segmented and armed as in *P. protothacae*. Outer edge of first segment of exopod of leg 1 without small spinules. Last segment of exopod of leg 4 with the formula II,I,5 in each of 10 females. Last segment of endopod of leg 4 (fig. 38) shaped as in previous species but shorter,  $57\mu \times 24\mu$ , its two terminal spines  $44\mu$  (outer) and  $60\mu$  (inner) in length.

Leg 5 (fig. 39) with the free segment  $43\mu \times 16\mu$ , ornamented with a small distal outer spinule, terminal seta  $50\mu$ , and subterminal spine  $18\mu$  with narrow lamellae.

Leg 6 as in *P. protothacae*.

Color in life in reflected light slightly amber, eye pale red, egg sacs whitish opaque.

MALE.—Body form much like that of male of *P. protothacae*, as shown in figures 24, 25. Prosome not tumid. Length (without the ramal setae) 1.38 mm (1.25–1.50 mm) and greatest width 0.44 mm (0.39–0.48 mm), based on 10 specimens measured in lactic acid.

Genital segment (fig. 40)  $260\mu \times 252\mu$ . Four postgenital segments  $69\mu \times 99\mu$ ,  $69\mu \times 94\mu$ ,  $51\mu \times 81\mu$ , and  $92\mu \times 79\mu$ , from anterior to posterior.

Caudal ramus like that of female,  $101\mu \times 30\mu$ .

Rostrum similar to that of female. First antenna  $297\mu$  long, segmented and armed as in male of *P. protothacae*. Second antenna as

in male of previous species. Labrum, mandible, paragnath, first maxilla, and second maxilla like those of the female. Maxilliped resembling that in *P. protothacae*, but claw  $150\mu$  along axis, and sclerotization of innermost seta on the second segment slightly different (fig. 41).

Postoral area as in female.

Legs 1-4 segmented and armed as in female. Last segment of endopod of leg 1 (fig. 42) with terminal spinules slightly coarser than in female.

Leg 5 (fig. 43) with the free segment  $32\mu \times 14\mu$ , without ornamentation, terminal seta  $55\mu$ , and subterminal spine  $10\mu$ , slender and setiform, without lamellae.

Leg 6 as in *P. protothacae*, the two setae  $31\mu$  and  $44\mu$  in length.

Spermatophore not seen.

Color in life in reflected light opaque white, eye pale red.

ETYMOLOGY.—The specific name *mesodesmatis* is formed from the generic name of the host.

COMPARISON WITH RELATED SPECIES.—*P. mesodesmatis* is very close to *P. protothacae*, but differs from it in several features. In *P. mesodesmatis* the prosome of the female is swollen, the ratio of length to width of the caudal ramus is 3.4:1 (instead of 4:1 as in *P. protothacae*), the first antennae are distinctly shorter, and certain other parts are relatively smaller as shown below.

Comparison of certain features of *Paranthessius protothacae* and *P. mesodesmatis* is as follows:

	<i>P. protothacae</i>	<i>P. mesodesmatis</i>
FEMALE		
Body size	1.62 x 0.54 mm	1.49 x 0.64 mm
Ratio of length to width of prosome	1.9:1	1.38:1
Nature of prosome	not tumid	tumid in nearly all specimens
Caudal ramus	$135\mu \times 34\mu$ (4:1)	$102\mu \times 30\mu$ (3.4:1)
Length of first antenna	$358\mu$	$285\mu$
Length of claw on second antenna	$77\mu$	$59\mu$
Formula for last segment of exopod of leg 4	usually II,I,5, but sometimes III,I,5	always II,I,5
Last segment of endopod of leg 4	$74\mu \times 28\mu$	$57\mu \times 24\mu$
Segment of leg 5 and two terminal elements	$52\mu \times 19\mu$ , seta $77\mu$ , spine $33\mu$	$43\mu \times 16\mu$ , seta $50\mu$ , spine $18\mu$
MALE		
Body size	1.44 x 0.44 mm	1.38 x 0.44 mm
Claw of maxilliped	$172\mu$	$150\mu$
Terminal spinules on last segment of endopod of leg 1	slender	coarse
Subterminal element of leg 5	$15\mu$ , spiniform	$10\mu$ , setiform

The differences observed between these two new species appear to be constant among the specimens studied, without overlap. The swollen prosome in the female of *P. mesodesmatis* occurred in both ovigerous and nonovigerous specimens. Bocquet and Stock (1959) stated that the prosome in *Paranthesius* tends to become larger and more swollen with the age of the females, doubtless because of the pressure exerted by the ovaries. In the preserved specimens of *P. mesodesmatis* it was impossible to separate the females by age since there was no way of being certain that egg sacs in some cases had not accidentally been broken off. The single less tumid mature female mentioned above may represent an individual that had not yet produced egg sacs.

## Literature Cited

BOCQUET, C., and STOCK, J. H.

1957. Copépodes parasites d'invertébrés des côtes de France, iv a-c: Le double parasitisme de *Sipunculus nudus* L. par *Myzomolgus stupendus* nov. gen., nov. sp., et *Catinia plana* nov. gen., nov. sp., copépodes cyclopoïdes très remarquables. Koninkl. Nederlandsch Akad. Wetensch., Amsterdam, Proc., ser. C, vol. 60, no. 3, pp. 410-431.
1958. Copépodes parasites d'invertébrés des côtes de France, vi: Description de *Paranthesius myxicolae* nov. sp., copépoide semi-parasite du Sabellidae *Myxicola infundibulum* (Rénier). Koninkl. Nederlandsch Akad. Wetensch., Amsterdam, Proc., ser. C, vol. 61, no. 2, pp. 243-253.
- 1958a. Copépodes parasites d'invertébrés des côtes de France, viii: Le genre *Ischnurella* Pelseneer, synonyme de *Paranthesius* Claus (Cyclopoida, Lichomolgidae). Koninkl. Nederlandsch Akad. Wetensch., Amsterdam, Proc., ser. C, vol. 61, no. 5, pp. 604-609.
1959. Copépodes parasites d'invertébrés des côtes de France, x: Sur les espèces de *Paranthesius* (Cyclopoida, Lichomolgidae) du groupe des *Herrmannella*, associées à des pélecypodes. Koninkl. Nederlandsch Akad. Wetensch., Amsterdam, Proc., ser. C, vol. 62, no. 3, pp. 238-249.
- 1959a. Copépodes parasites d'invertébrés des côtes de la Manche, vi: Redescription de *Paranthesius anemoniae* Claus (Copepoda Cyclopoida) parasite d'*Anemonia sulcata* (Pennant). Arch. Zool. Exp. Gén., vol. 98, notes et revue, no. 1, pp. 43-53.

BRIAN, A.

1924. Parasitologia Mauritanica: Matériaux pour la faune parasitologique en Mauritanie: Arthropoda (1<sup>re</sup> partie) Copepoda. Bull. Comité d'Études Hist. Sci. Afrique Occidentale Française, 1924 (July-September) pp. 4-66.

CANU, E.

1891. Les copépodes marins du Boulonnais, v: Les semi-parasites. Bull. Sci. France et Belgique, vol. 23, pp. 467-487.

CLAUS, C.

1889. Über neue oder wenig bekannte halbparasitische Copepoden, insbesondere die Lichomolgiden- und Ascomyzontiden-Gruppe. Arb. Zool. Inst. Univ. Wien, vol. 8, pp. 327-370.

EDWARDS, C. L.

1891. Beschreibung einiger neuen Copepoden und eines neuen copepodenähnlichen Krebses, *Leuckartella paradoxa*. Inaugural dissertation, University of Leipzig, pp. 1-33. Berlin.

GOTTO, R. V.

1961. A new lichomolgid copepod, and the occurrence of two little-known cyclopoids in British waters. Crustaceana, vol. 3, no. 2, pp. 85-92.

HEEGAARD, P.

1944. A new copepod (*Scambicornus hamatus*) parasitic on a Japanese holothurian. Vidensk. Medd. Dansk Naturh. Foren., vol. 107, pp. 359-366.

## ILLG, P. L.

1949. A review of the copepod genus *Paranthessius* Claus. Proc. U.S. Nat. Mus., vol. 99, no. 3245, pp. 391-428.

## KOSSMANN, R.

1877. Entomostraca (1. Theil: Lichomolgidae). Vol. 4 in Zoologische Ergebnisse . . . Reise in die Küstengebiete des Rothen Meeres, pp. 1-24.

## MONOD, T., and DOLLFUS, R.-PH.

1932. Les copépodes parasites de mollusques. Ann. Parasitol. Hum. Comp., vol. 10, pp. 129-204.

## NICHOLLS, A. G.

1944. Littoral Copepoda from South Australia (II): Calanoida, Cyclopoida, Notodelphyoida, Monstrilloida and Caligoida. Rec. South Australian Mus., vol. 8, no. 1, pp. 1-62.

## NORMAN, A. M., and Scott, T.

1905. Crustacea Copepoda new to science from Devon and Cornwall. Ann. Mag. Nat. Hist., ser. 7, vol. 15, pp. 284-300.

## PELSENEER, P.

1929. Copépodes parasites de mollusques. Ann. Soc. Roy. Zool. Belgique, 1928, vol. 59, pp. 33-49.

## PESTA, O.

1909. Beiträge zur Kenntnis parasitischer Copepoden. Denkschr. Kais. Akad. Wiss., Wien, Math.-naturwiss. Kl., vol. 84, pp. 257-267.

## REDDIAH, K.

1960. A new lichomolgid copepod, *Paranthessius colmani* n. sp., from the *Laevicardium crassum* (Gmelin) Lamellibranchiata-Cardiidae. Ann. Mag. Nat. Hist., 1959, ser. 13, vol. 2, no. 19, pp. 439-443.

## REDDIAH, K., and WILLIAMSON, D. I.

1959. On *Modiolicola inermis* Canu and *Modiolicola maxima* (Thompson), lichomolgid copepods associated with pectinid lamellibranchs. Ann. Mag. Nat. Hist., 1958, vol. 13, no. 1, pp. 689-701.

## SARS, G. O.

1918. An account of the Crustacea of Norway with short descriptions and figures of all the species, vol. 6: Copepoda Cyclopoida, parts 13 and 14: Lichomolgidae (concluded), Oncaeidae, Corycaeidae, Ergasilidae, Clausiidae, Eunicicolidae, Supplement, pp. 173-225. Bergen: Bergen Museum.

## SCOTT, T.

1903. Notes on some Copepoda from the Arctic Seas collected in 1890 by the Rev. Canon A. M. Norman, F.R.S. Ann. Mag. Nat. Hist., ser. 7, vol. 11, no. 61, pp. 4-32.

## SEWELL, R. B. S.

1949. The littoral and semi-parasitic Cyclopoida, the Monstrilloida and Notodelphyoida. In John Murray Expedition 1933-34, Scientific Reports, vol. 9, no. 2, pp. 17-199.

## STOCK, J. H.


1964. On *Scambicornus* Heegard, 1944, a senior synonym of *Preherrmannella* Seymour Sewell, 1949 (Copepoda, Cyclopoida). Beaufortia, vol. 10, no. 123, pp. 183-192.

**THOMPSON, I. C.**


1897. Appendix: Note on dredging and tow netting in Puget Sound, Pacific Coast. *In* Herdman, W. A., Thompson, I. C., and Scott, A., On the plankton collected continuously during two traverses of the North Atlantic in the summer of 1897; with descriptions of new species of Copepoda; and an appendix on dredging in Puget Sound. *Trans. Liverpool Biol. Soc.*, vol. 12, pp. 33-90.

**THOMPSON, I. C., and SCOTT, A.**


1903. Report on the Copepoda collected by Professor Herdman, at Ceylon, in 1902. *In* Herdman, Report to the Government of Ceylon on the Pearl Oyster Fisheries of the Gulf of Manaar, pt. I, Suppl. Rept. no. 7, pp. 227-307.




FIGURES 1-7.—*Paranthessius protothacae*, new species, female: 1, body, dorsal (A); 2, body, lateral (A); 3, urosome, dorsal (B); 4, segments of legs 4 and 5, ventral (C); 5, segments of legs 4 and 5, lateral (C); 6, area of attachment of egg sac, dorsal (D); 7, caudal ramus, dorsal (E).



FIGURES 8-15.—*Paranthessius protothacae*, new species, female: 8, rostrum, ventral (E); 9, outline of rostrum, lateral (E); 10, first antenna, dorsal (E); 11, third segment of first antenna, ventral (F); 12, second antenna, posterior (E); 13, edge of labrum, paragnaths, and lingua, ventral (E); 14, mandible, posterior (F); 15, first maxilla, anterior (F).



FIGURES 16-20.—*Paranthessius protothacae*, new species, female: 16, second maxilla, inner (F); 17, maxilliped, antero-inner (F); 18, oral and postoral area, ventral (B); 19, leg 1, anterior (E); 20, leg 2, anterior (E).



FIGURES 34-43.—*Paranthessius mesodesmatis*, new species, female: 34, body, dorsal (A); 35 body, lateral (A); 36, urosome, dorsal (B); 37, outline of rostrum, lateral (E); 38, endopod of leg 4, anterior (E); 39, leg 5, dorsal (F). Male: 40, urosome, dorsal (B); 41, innermost seta on second segment of maxilliped, inner (G); 42, last segment of endopod of leg 1, anterior (F); 43, leg 5, dorsal (F).







Humes, Arthur G. 1967. "Cyclopoid copepods of the genus *Paranthesius* associated with marine pelecypods in Chile." *Proceedings of the United States National Museum* 124(3628), 1–18.

<https://doi.org/10.5479/si.00963801.124-3628.1>.

**View This Item Online:** <https://www.biodiversitylibrary.org/item/32359>

**DOI:** <https://doi.org/10.5479/si.00963801.124-3628.1>

**Permalink:** <https://www.biodiversitylibrary.org/partpdf/10186>

**Holding Institution**

Smithsonian Libraries

**Sponsored by**

Smithsonian

**Copyright & Reuse**

Copyright Status: NOT\_IN\_COPYRIGHT

Rights: <https://www.biodiversitylibrary.org/permissions/>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.